



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE.—The following equations (g) and (h), and concluding remark, was enclosed by Prof. Philbrick with the corrected “proof” of his paper, pp. 9–14, and should have appeared in connection with that paper; but by accident the proof was not rec’d till after the sheet was printed, hence their appearance here, and the corrections of the proof, in the *Errata* on p. 32.

“Also

$$\int \frac{\sin^m x \cos^r x dx}{(a+b \sec x)^n} = \int \frac{\sin^m x \cos^{r+n} x dx}{(b+a \cos x)^n}, \quad (g)$$

which may be integrated by (a) or by (b); and

$$\int \frac{\sin^m x \cos^r x dx}{a+b \coth x)^n} = \int \frac{\sin^{m+n} x \cos^r x dx}{(b+a \sin x)^n}. \quad (h)$$

which may be integrated by (c) or (d).

Probably almost any combination of trigonometric functions may be integrated, directly or by transformation, by the general formulas above, or by others easily derived from them.”

CALCULATION OF TRANSIT OF VENUS BY PROF. BARBOUR.—This calculation is made, by T. H. Safford, Jr.’s modification of Bessel’s method, for the position of Louisville, Ky., N. Lat. $38^\circ 14' 57''.78$, Long. $85^\circ 45' 52''.53$ W.; transit to occur on Dec. 6th 1882.

The formulæ will be found in Chauvenet’s *Spher. and Pract. Ast.*, Vol. I, and in the *Amer. Naut. Almanac* for 1882.

$$a = A - h \sin(\mu - \lambda), \quad b = B - EK + Gh \cos(\mu - \lambda),$$

$c = -C + FK - Hh \cos(\mu - \lambda)$, $m = \sqrt{bc}$. If $m = a$ the assumed time is correct.—To find time of 2nd contact. Put φ' = geocentric latitude of the place; λ = longitude West from Greenwich; ρ = dist. from Earth’s center; $h = \rho \cos \varphi'$, and $K = \rho \sin \varphi'$.

An easy method of calculating h and K is provided by *Amer. Naut. Almanac* for 1882, p. 499. $\rho \cos \varphi' = F' \cos \varphi$, and $\rho \sin \varphi' = G' \sin \varphi$, in which φ = geographical latitude, and logs F' and G' are given in a table.

$$\begin{array}{ll} \log \cos \varphi = 9.8950487 & \log \sin \varphi = 9.7917506 \\ \log F' = .000578 & \log G' = n .0024355 \\ \log h = 9.8956267 & \log K = n 9.7941861 \end{array}$$

(n before a log. means that the number corresponding is negative.)

To find a . $\mu = 38^\circ 13' 3''$ for epoch $2^h 24^m 57^s$ (Gr. mean time).

$$\begin{array}{l} \lambda = 85^\circ 45' 52''.53; \therefore \mu - \lambda = -47^\circ 32' 49''.53 \text{ or } 312^\circ \\ 27' 10''.47. \therefore \log \sin(\mu - \lambda) = n 9.8679576; \log \cos(\mu - \lambda) = 9.8292395 \\ \log h = 9.8956267 \end{array}$$

$$\log h + \log \sin(\mu - \lambda) = 9.7635843. \therefore h \sin(\mu - \lambda) = .580209.$$

$$A, \text{ for given epoch, } = 18.8281658; +.580209 = 19.4083748 = a.$$